



Mac Lane's coherence theorem expressed as a word problem

Paul-André Melliès

► To cite this version:

| Paul-André Melliès. Mac Lane's coherence theorem expressed as a word problem. 2003. hal-00154213

HAL Id: hal-00154213

<https://hal.science/hal-00154213>

Preprint submitted on 19 Jun 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MacLane's coherence theorem expressed as a word problem

Paul-André Mellies
Preuves, Programmes, Systèmes
CNRS UMR-7126, Université Paris 7
<mellies@pps.jussieu.fr>

In this draft manuscript, we reduce the coherence theorem for braided monoidal categories to the resolution of a word problem, and the construction of a category of fractions. The technique explicates the combinatorial nature of that particular coherence theorem.

1. Introduction

Let \mathcal{B} denote the category of braids and \mathcal{M} any braided monoidal category. Let $\text{Br}(\mathcal{B}, \mathcal{M})$ denote the category of *strong* braided monoidal functors from \mathcal{B} to \mathcal{M} and monoidal natural transformations between them. All definitions are recalled in section 5. The coherence theorem for braided monoidal categories is usually stated as follows:

Theorem 1. The categories $\text{Br}(\mathcal{B}, \mathcal{M})$ and \mathcal{M} are equivalent.

We prove the coherence theorem in four independent steps.

To that purpose, we introduce the category \mathcal{A}

- whose objects are binary trees of \otimes , with leaves 0 and 1,
- whose morphisms are sequences of rewriting steps $\alpha(a, b, c), \lambda(a), \rho(a), \gamma(a, b), \alpha^{-1}(a, b, c), \lambda^{-1}(a), \rho^{-1}(a), \gamma^{-1}(a, b)$, quotiented by the laws of braided monoidal categories.

1.1. First step

Consider a braided monoidal category \mathcal{M} , and the category $\text{SBr}(\mathcal{A}, \mathcal{M})$ of *strict* braided monoidal functors and monoidal natural transformations from \mathcal{A} to \mathcal{M} . We prove that \mathcal{M} and $\text{SBr}(\mathcal{A}, \mathcal{M})$ are isomorphic. More precisely, we prove that the functor

$$[-] : \text{SBr}(\mathcal{A}, \mathcal{M}) \longrightarrow \mathcal{M}$$

which associates to a strict natural transformation

$$\theta : (F, F_2, F_0) \longrightarrow (G, G_2, G_0)$$

the morphism

$$\theta_1 : F(1) \longrightarrow G(1)$$

in \mathcal{M} , is reversible.

To every object X in the category \mathcal{M} , we associate a strict braided monoidal functor $\llbracket X \rrbracket$ defined as follows:

$$\llbracket X \rrbracket(0) = e \quad \llbracket X \rrbracket(1) = X \quad \llbracket X \rrbracket(a \otimes b) = \llbracket X \rrbracket(a) \otimes \llbracket X \rrbracket(b)$$

and

$$\llbracket X \rrbracket(\alpha) = \alpha \quad \llbracket X \rrbracket(\lambda) = \lambda \quad \llbracket X \rrbracket(\rho) = \rho \quad \llbracket X \rrbracket(\gamma) = \gamma$$

This defines a functor because the image of a commutative diagrams in \mathcal{A} is always commutative in \mathcal{M} . The functor is strict braided monoidal by construction.

To every morphism $f : X \rightarrow Y$ in the category \mathcal{M} , we associate a family of morphisms $\llbracket f \rrbracket$ indexed by objects of \mathcal{A} , as follows:

$$\llbracket f \rrbracket_0 = \text{id}_e \quad \llbracket f \rrbracket_1 = f \quad \llbracket f \rrbracket_{a \otimes b} = \llbracket f \rrbracket_a \otimes \llbracket f \rrbracket_b$$

The family defines a natural transformation $\llbracket f \rrbracket : \llbracket X \rrbracket \rightarrow \llbracket Y \rrbracket$ because the maps α, λ, ρ , and γ are supposed natural in \mathcal{M} . The natural transformation $\llbracket f \rrbracket$ is monoidal by construction.

The map $\llbracket - \rrbracket$ is functorial from \mathcal{M} to $\text{SBr}(\mathcal{A}, \mathcal{M})$ and it is easy to see that it defines the inverse of the evaluation functor $[-] : \text{SBr}(\mathcal{A}, \mathcal{M}) \rightarrow \mathcal{M}$. We conclude.

1.2. Second step

We introduce the notion of *contractible* category. A category Ξ is contractible when:

- there exists at most one morphism between two objects of Ξ ,
- every morphism of Ξ is reversible.

In other words, a category Ξ is contractible when it is a preorder category and a groupoid.

Consider a contractible subcategory Ξ of a category \mathcal{C} , full on objects. Write $a \simeq_{\Xi}^{\text{obj}} b$ for two objects a and b of \mathcal{C} , when there exists a morphism $a \rightarrow b$ in Ξ . Write $f \simeq_{\Xi}^{\text{map}} g$ for two morphisms $f : a \rightarrow b$ and $g : c \rightarrow d$ when there exists two morphisms $a \rightarrow c$ and $b \rightarrow d$ in Ξ making the following diagram commute:

$$\begin{array}{ccc} a & \xrightarrow{f} & b \\ \downarrow & & \downarrow \\ c & \xrightarrow{g} & d \end{array}$$

The relations $\simeq_{\Xi}^{\text{obj}}$ and $\simeq_{\Xi}^{\text{map}}$ are equivalence relations on objects and morphisms of \mathcal{C} , respectively. We call *orbit* of an object a or morphism f its equivalence class wrt. $\simeq_{\Xi}^{\text{obj}}$ or $\simeq_{\Xi}^{\text{map}}$. The quotient category \mathcal{C}/Ξ is defined as follows:

- its objects are the orbits of objects,
- its morphisms $M \rightarrow N$ are the orbits of maps,
- its identities and composition are induced from \mathcal{C} .

The two categories \mathcal{C} and \mathcal{C}/Ξ are equivalent. Indeed, there exists a “projection” functor

$$F : \mathcal{C} \rightarrow \mathcal{C}/\Xi$$

which maps every morphism $f : M \rightarrow N$ to its orbit, and an “embedding” functor

$$G : \mathcal{C}/\Xi \rightarrow \mathcal{C}$$

depending on the choice, for every orbit wrt. $\simeq_{\Xi}^{\text{obj}}$, of an object in that class. Clearly, $FG = \text{id}_{\mathcal{C}/\Xi}$, and there exists a natural transformation $GF \rightarrow \text{id}_{\mathcal{C}}$.

1.3. Third step

Consider a braided monoidal category \mathcal{M} , and the hom-category $\text{Br}(\mathcal{A}, \mathcal{M})$ of strong braided monoidal functors and monoidal natural transformations from \mathcal{A} to \mathcal{M} . We prove that the categories $\text{Br}(\mathcal{A}, \mathcal{M})$ and $\text{SBr}(\mathcal{A}, \mathcal{M})$ are equivalent.

Define Ξ as the subcategory of $\text{Br}(\mathcal{A}, \mathcal{M})$ containing all reversible maps $\theta : (F, F_2, F_0) \xrightarrow{\sim} (G, G_2, G_0)$ such that $\theta_1 = \text{id}_{F(1)}$. Every identity of $\text{Br}(\mathcal{A}, \mathcal{M})$ is element of Ξ . The main observation is that the category Ξ is contractible. Indeed, once the equality $\theta_1 = \text{id}_{F(1)}$ is fixed, the component morphisms θ_0 and $\theta_{a \otimes b}$ of the natural transformation θ are uniquely determined by the commutative diagrams

$$\begin{array}{ccc} F(a) \otimes F(b) & \xrightarrow{F_2(a, b)} & F(a \otimes b) \\ \theta_a \otimes \theta_b \downarrow & & \downarrow \theta_{a \otimes b} \\ G(a) \otimes G(b) & \xrightarrow{G_2(a, b)} & G(a \otimes b) \end{array} \quad \begin{array}{ccc} e & \xrightarrow{F_0} & F(0) \\ \parallel & & \downarrow \theta_0 \\ e & \xrightarrow{G_0} & G(0) \end{array}$$

Moreover, every identity of $\text{Br}(\mathcal{A}, \mathcal{M})$ is element of Ξ . We may therefore consider the category $\text{Br}(\mathcal{A}, \mathcal{M})/\Xi$, which we know is equivalent to $\text{Br}(\mathcal{A}, \mathcal{M})$.

We prove that the categories $\text{Br}(\mathcal{A}, \mathcal{M})/\Xi$ and $\text{SBr}(\mathcal{A}, \mathcal{M})$ are isomorphic. We need to prove that for every strong monoidal functor (G, G_2, G_0) from \mathcal{B} to \mathcal{M} , there exists a strict monoidal functor F , and a map $\theta : F \xrightarrow{\sim} (G, G_2, G_0)$ in Ξ . Consider a family of isomorphisms $\theta_a : F_a \xrightarrow{\sim} G_a$

$$\theta_0 = G_0 \quad \theta_1 = \text{id}_e \quad \theta_{a \otimes b} = G_2(a, b) \circ (\theta_a \otimes \theta_b)$$

indexed by objects of \mathcal{A} . Then, associate to every morphism $f : a \rightarrow b$ in \mathcal{A} , the morphism

$$F(f) = \theta_b^{-1} \circ G(f) \circ \theta_a$$

in \mathcal{M} . A close look at the diagram shows that this defines a strict braided monoidal functor $F : \mathcal{A} \rightarrow \mathcal{M}$ and a monoidal natural transformation $\theta : F \xrightarrow{\sim} G$ in Ξ . We conclude.

1.4. Fourth step

After steps 1. 2. and 3. the proof of coherence reduces to the comparison of two “free categories”:

- the “formal” braided monoidal category \mathcal{A} of \otimes -trees and rewriting paths, quotiented by the commutativities of braided monoidal categories,
- the “geometric” braided monoidal category \mathcal{B} of braids.

This is the most interesting and difficult part of the proof, in fact the first and only time combinatorics plays a role. We prove

1. that the subcategory of \mathcal{A} consisting only of α , ρ and λ maps, and their inverses, is contractible,
2. that the quotient category is isomorphic to the braid category \mathcal{B} , and equivalent to \mathcal{A} through strong braided monoidal functors.

The first and second assertions are established in sections 3 and 4 respectively.

1.5. Conclusion

The categories $\text{Br}(\mathcal{A}, \mathcal{M})$ and \mathcal{M} are equivalent by steps 1. 2. and 3. and the categories $\text{Br}(\mathcal{A}, \mathcal{M})$ and $\text{Br}(\mathcal{B}, \mathcal{M})$ are equivalent by step 4. This proves theorem 1.

2. A word problem

Let Σ be the category

- whose objects are \otimes -trees with leaves 0 and 1,
- whose morphisms are sequences of $\alpha(a, b, c)$, $\lambda(a)$ and $\rho(a)$, quotiented by the monoidal law (1) the commutativity laws (5) (6) and the additional commutativity triangles (2).

$$\begin{array}{ccc} a \otimes b & \xrightarrow{f \otimes b} & a' \otimes b \\ a \otimes g \downarrow & & \downarrow a' \otimes g \\ a \otimes b' & \xrightarrow{f \otimes b'} & a' \otimes b' \end{array} \quad \text{for} \quad \begin{array}{ccc} a & \xrightarrow{f} & a' \\ b & \xrightarrow{g} & b' \end{array} \quad (1)$$

$$\begin{array}{ccc} e \otimes (a \otimes b) & \xrightarrow{\alpha} & (e \otimes a) \otimes b \\ \lambda \downarrow & & \downarrow \lambda \otimes b \\ a \otimes b & \xlongequal{\quad} & a \otimes b \end{array} \quad \begin{array}{ccc} a \otimes (b \otimes e) & \xrightarrow{\alpha} & (a \otimes b) \otimes e \\ a \otimes \rho \downarrow & & \downarrow \rho \\ a \otimes b & \xlongequal{\quad} & a \otimes b \end{array} \quad (2)$$

Theorem 2. The category Σ verifies the three following properties:

- the category has pushouts,
- every morphism is epi,
- from every object a , there exists a morphism $a \longrightarrow b$ to a normal object b .

Proof. The proof appears in (Melliès, 2002). It is motivated by works of rewriters like Lévy (Lévy, 1978; Huet and Lévy, 1979) and algebraists like Dehornoy (Dehornoy, 1998). \square

Definition 1. An object a in a category \mathcal{C} is called normal when id_a is the only morphism outgoing from a .

3. Calculus of fractions

3.1. Definition

For every category \mathcal{C} and class Σ of morphisms in \mathcal{C} , there exists a universal solution to the problem of “reversing” the morphisms in Σ . More explicitly, there exists a category $\mathcal{C}[\Sigma^{-1}]$ and a functor

$$P_\Sigma : \mathcal{C} \longrightarrow \mathcal{C}[\Sigma^{-1}]$$

such that:

- the functor P_Σ maps every morphism of Σ to reversible morphism,
- if a functor $F : \mathcal{C} \longrightarrow \mathcal{M}$ makes every morphism of Σ reversible, then F factors as $F = G \circ P_\Sigma$ for a unique functor $G : \mathcal{C}[\Sigma^{-1}] \longrightarrow \mathcal{M}$.

The class Σ is a **calculus of left fraction** in the category \mathcal{C} , see (Gabriel and Zisman, 1967), when

- Σ contains the identities of \mathcal{C} ,
- Σ is closed under composition,
- each diagram $Y \xleftarrow{s} X \xrightarrow{f} Z$ where $s \in \Sigma$ may be completed into a commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ s \downarrow & & \downarrow t \\ Z & \xrightarrow{g} & X' \end{array} \quad \text{where } t \in \Sigma$$

- each commutative diagram

$$X' \xrightarrow{s} X \xrightarrow[g]{f} Y \quad \text{where } s \in \Sigma$$

may be completed into a commutative diagram

$$X \xrightarrow[g]{f} Y \xrightarrow[t]{g} Y' \quad \text{where } t \in \Sigma$$

The property of left fraction calculus enables an elegant definition of the category $\mathcal{C}[\Sigma^{-1}]$ and functor $P_\Sigma : \mathcal{C} \rightarrow \mathcal{C}[\Sigma^{-1}]$. The category $\mathcal{C}[\Sigma^{-1}]$ is defined as follows.

Its objects are the objects of \mathcal{C} , and its morphisms $X \rightarrow Y$ are the equivalence classes of pairs (f, s) of morphisms of \mathcal{C}

$$X \xrightarrow{f} \cdot \xleftarrow{s} Y \quad \text{where } s \in \Sigma$$

under the equivalence relation which identifies two pairs $X \xrightarrow{f} Z \xleftarrow{s} Y$ and $X \xrightarrow{g} Z' \xleftarrow{t} Y$ when there exists a pair $X \xrightarrow{h} Z'' \xleftarrow{u} Y$ and two morphisms $Z \xrightarrow{f'} Z'' \xleftarrow{g'} Z'$ forming a commutative diagram


$$\begin{array}{ccccc} & & Z & & \\ & f \nearrow & \downarrow f' & \nwarrow s & \\ X & \xrightarrow{h} & Z'' & \xleftarrow{u} & Y \\ & g \searrow & \uparrow g' & \nearrow t & \\ & & Z' & & \end{array} \quad \text{where } u \in \Sigma$$

3.2. Application

By theorem 2, the class Σ is a calculus of fraction in the category Σ . But we can prove a bit more. A consequence of the pushout property, and definition of a normal object, is that any two morphisms $a \rightarrow e$ and $b \rightarrow e$ to a normal object e are equal in Σ . So, consider two morphisms in $\Sigma[\Sigma^{-1}]$, represented as pairs in Σ

$$a \xrightarrow{f} c \xleftarrow{s} b \quad a \xrightarrow{g} d \xleftarrow{t} b$$

Any two morphisms $b \rightarrow e$ and $c \rightarrow e$ to a normal object e in Σ , make the following diagram commute in Σ :



We obtain

Theorem 3. The category $\Sigma[\Sigma^{-1}]$ is contractible.

4. From word problem to coherence theorem

In section 3 we have proved that the category $\Sigma[\Sigma^{-1}]$ consisting of $\alpha(a, b, c)$, $\lambda(a)$ $\rho(a)$ steps and their inverse, is contractible.

- In step 1. we prove that $\Sigma[\Sigma^{-1}]$ is a subcategory of \mathcal{A} , full on objects.
- In step 2. we prove that the category $\mathcal{A}/\Sigma[\Sigma^{-1}]$ is braided monoidal, and embeds in \mathcal{A} through strong braided monoidal functors,
- in step 3. we prove that the category $\mathcal{A}/\Sigma[\Sigma^{-1}]$ is isomorphic to the usual braid category \mathcal{B} .

4.1. First step

The two triangles (2) are commutative in \mathcal{A} , as in every monoidal category, see exercise VII.§1.1. in (Mac Lane, 1991). This ensures the existence of a functor $\Sigma \rightarrow \mathcal{A}$. By definition, this functor induces a functor $\Sigma[\Sigma^{-1}] \rightarrow \mathcal{A}$. This functor is injective on objects. It is also faithful for the obvious reason that $\Sigma[\Sigma^{-1}]$ is contractible. Thus, $\Sigma[\Sigma^{-1}] \rightarrow \mathcal{A}$ defines an embedding of the contractible category $\Sigma[\Sigma^{-1}]$ into \mathcal{A} . The embedding is full on objects. This enables to consider the quotient category $\mathcal{A}/\Sigma[\Sigma^{-1}]$.

4.2. Second step

The category $\mathcal{A}/\Sigma[\Sigma^{-1}]$ becomes strict monoidal with the following definition. Let $f : a \rightarrow b$ and $g : c \rightarrow d$ be two morphisms of $\mathcal{A}/\Sigma[\Sigma^{-1}]$ represented by two morphisms $f_0 : a_0 \rightarrow b_0$ and $g_0 : c_0 \rightarrow d_0$ in the category \mathcal{A} . Define the tensor product $f \otimes g : a \otimes b \rightarrow c \otimes d$ as the projection of the tensor product $f_0 \otimes g_0 : a_0 \otimes c_0 \rightarrow b_0 \otimes d_0$ and the unit object e as the orbit of the unit in \mathcal{A} . Correctness follows from monoidality of \otimes in \mathcal{A} . Monoidality of \otimes is not difficult to prove, and diagrams (5) (6) are obvious.

The category $\mathcal{A}/\Sigma[\Sigma^{-1}]$ is braided monoidal. Let a and b be two objects of the quotient category $\mathcal{A}/\Sigma[\Sigma^{-1}]$ represented by two objects a_0 and b_0 in the category \mathcal{A} . Define the braiding $\gamma(a, b) : a \otimes b \rightarrow b \otimes a$ in the quotient category as the orbit of the braiding

$\gamma(a_0, b_0) : a_0 \otimes b_0$. Correctness of the definition and naturality of γ follow from naturality of γ in \mathcal{A} . Diagrams (7) (8) (9) follow from the definition of γ in $\mathcal{A}/\Sigma[\Sigma^{-1}]$.

Moreover,

- the “projection” functor $F : \mathcal{A} \rightarrow \mathcal{A}/\Sigma[\Sigma^{-1}]$ is strict braided monoidal. Diagrams (10) (11) are trivial, and diagram (12) follows from the definition of the braiding γ in $\mathcal{A}/\Sigma[\Sigma^{-1}]$.
- the “embedding” functor $(G, G_2, G_0) : \mathcal{B}/\Xi \rightarrow \mathcal{A}$ is strong braided monoidal, with the morphisms $G_0 : e \rightarrow G(e')$ and $G_2(a, b) : G(a) \otimes G(b) \rightarrow G(a \otimes b)$ determined as morphisms of Σ . Diagrams (10) (11) hold because every maps are in $\Sigma[\Sigma^{-1}]$, and diagram (12) holds because of the definition of the braiding in $\mathcal{A}/\Sigma[\Sigma^{-1}]$.

4.3. Third step

We prove that $\mathcal{A}/\Sigma[\Sigma^{-1}]$ and \mathcal{B} are isomorphic categories. The objects of $\mathcal{A}/\Sigma[\Sigma^{-1}]$ are the natural numbers. The morphisms of $\mathcal{A}/\Sigma[\Sigma^{-1}]$ are sequences of $\rho(m, n)$ and $\rho^{-1}(m, n)$ steps, modulo monoidality, and commutativity of the triangles induced by (8) (9):

$$\begin{array}{ccc}
 m \otimes n \otimes p & \xrightarrow{\gamma(m \otimes n, p)} & p \otimes m \otimes n \\
 \downarrow m \otimes \gamma(n, p) & & \parallel \\
 m \otimes p \otimes n & \xrightarrow{\gamma(m, p) \otimes n} & p \otimes m \otimes n
 \end{array}
 \quad
 \begin{array}{ccc}
 m \otimes n \otimes p & \xrightarrow{\gamma(m, n \otimes p)} & n \otimes p \otimes m \\
 \downarrow \gamma(m, n) \otimes p & & \parallel \\
 n \otimes m \otimes p & \xrightarrow{n \otimes \gamma(m, p)} & n \otimes p \otimes m
 \end{array}
 \quad (3)$$

We construct a strict monoidal functor from $\Sigma[\Sigma^{-1}]$ to \mathcal{B} by interpreting each $\gamma(m, n)$ by the braiding $\gamma_{\mathcal{B}}(m, n) : m + n \rightarrow n + m$ in \mathcal{B} consisting in permuting n braids over m braids. The functor is full. The only difficult point to prove is that it is faithful.

This reduces to proving that the hexagonal diagram generating equality of “braids” in \mathcal{B} commutes (already) in the category $\mathcal{A}/\Sigma[\Sigma^{-1}]$:

$$\begin{array}{ccccc}
 & & m \otimes n \otimes p & & \\
 & \swarrow \gamma(m, n) \otimes p & & \searrow m \otimes \gamma(n, p) & \\
 n \otimes m \otimes p & & & & m \otimes p \otimes n \\
 \downarrow n \otimes \gamma(m, p) & & & & \downarrow \gamma(m, p) \otimes n \\
 n \otimes p \otimes m & & & & p \otimes m \otimes n \\
 & \swarrow \gamma(n, p) \otimes m & & \swarrow p \otimes \gamma(m, n) & \\
 & & p \otimes n \otimes m & &
 \end{array}
 \quad (4)$$

The diagram is a nice “geometric” consequence of the commutative triangles 3, and monoidality of $\mathcal{A}/\Sigma[\Sigma^{-1}]$.

We conclude that $\mathcal{A}/\Sigma[\Sigma^{-1}]$ and the category \mathcal{B} of braids are isomorphic.

References

P. Dehornoy. *On completeness of word reversing*. Discrete mathematics, to appear.

- P. Gabriel and M. Zisman.** *Calculus of Fractions and Homotopy Theory*. Ergebnisse des Mathematik. Springer-Verlag, 1967.
- G. Huet, J.-J. Lévy.** *Call by Need Computations in Non-Ambiguous Linear Term Rewriting Systems*. Rapport de recherche INRIA 359, 1979. Reprinted as: Computations in orthogonal rewriting systems. In J.-L. Lassez and G. D. Plotkin, editors, *Computational Logic; Essays in Honor of Alan Robinson*, pages 394–443. MIT Press, 1991.
- A. Joyal and R. Street.** Braided tensor categories. *Adv. Math.* 102, 20-78, 1993.
- C. Kassel.** *Quantum groups*. Graduate Texts in Mathematics, Springer-Verlag, 1991.
- J.-J. Lévy.** *Réductions correctes et optimales dans le λ -calcul*. Thèse de Doctorat d'Etat, Université Paris VII, 1978.
- S. Mac Lane.** *Categories for the working mathematician*. Second edition. Graduate Texts in Mathematics, Springer-Verlag, 1991.
- P-A. Mellies.** Axiomatic Rewriting Theory 6: Residual Theory Revisited. Proceedings of RTA 2002.

5. Appendix: definitions of braided monoidal categories, functors and natural transformations

5.1. Monoidal category

A monoidal category \mathcal{M} is a category with a bifunctor

$$\otimes : \mathcal{M} \times \mathcal{M} \longrightarrow \mathcal{M}$$

which is associative up to a natural isomorphism

$$\alpha : a \otimes (b \otimes c) \longrightarrow (a \otimes b) \otimes c$$

equipped with an element e , which is a unit up to natural isomorphisms

$$\lambda : e \otimes a \longrightarrow a \quad \rho : a \otimes e \longrightarrow a$$

These maps must make Mac Lane's famous "pentagon" commute

$$\begin{array}{ccc} a \otimes (b \otimes (c \otimes d)) & \xrightarrow{\alpha} & (a \otimes b) \otimes (c \otimes d) \\ \downarrow a \otimes \alpha & & \downarrow \alpha \\ a \otimes ((b \otimes c) \otimes d) & \xrightarrow{\alpha} (a \otimes (b \otimes c)) \otimes d \xrightarrow{\alpha \otimes d} & ((a \otimes b) \otimes c) \otimes d \end{array} \quad (5)$$

as well as the triangle:

$$\begin{array}{ccc} a \otimes (e \otimes b) & \xrightarrow{\alpha} & (a \otimes e) \otimes b \\ \downarrow a \otimes \lambda & & \downarrow \rho \otimes b \\ a \otimes b & \xlongequal{\quad} & a \otimes b \end{array} \quad (6)$$

A monoidal category is *strict* when all α , λ and ρ are identities.

5.2. Braided monoidal category

A braiding for a monoidal category \mathcal{M} consists of a family of isomorphisms

$$\gamma_{a,b} : a \otimes b \longrightarrow b \otimes a$$

natural in a and b , which satisfy the commutativity

$$\begin{array}{ccc} a \otimes e & \xrightarrow{\gamma} & e \otimes a \\ \rho \downarrow & & \downarrow \lambda \\ a & \xlongequal{\quad} & a \end{array} \quad (7)$$

and makes both following hexagons commute:

$$\begin{array}{ccccc} a \otimes (b \otimes c) & \xrightarrow{\alpha} & (a \otimes b) \otimes c & \xrightarrow{\gamma} & c \otimes (a \otimes b) \\ \downarrow a \otimes \gamma & & & & \downarrow \alpha \\ a \otimes (c \otimes b) & \xrightarrow{\alpha} & (a \otimes c) \otimes b & \xrightarrow{\gamma \otimes b} & (c \otimes a) \otimes b \end{array} \quad (8)$$

$$\begin{array}{ccccc} (a \otimes b) \otimes c & \xrightarrow{\alpha^{-1}} & a \otimes (b \otimes c) & \xrightarrow{\gamma} & (b \otimes c) \otimes a \\ \downarrow \gamma \otimes c & & & & \downarrow \alpha^{-1} \\ (b \otimes a) \otimes c & \xrightarrow{\alpha^{-1}} & b \otimes (a \otimes c) & \xrightarrow{b \otimes \gamma} & b \otimes (c \otimes a) \end{array} \quad (9)$$

5.3. Monoidal functor

A monoidal functor $(F, F_2, F_0) : \mathcal{M} \rightarrow \mathcal{M}'$ between monoidal categories \mathcal{M} and \mathcal{M}' :

- an ordinary functor $F : \mathcal{M} \rightarrow \mathcal{M}'$,
- for objects a, b in \mathcal{M} morphisms in \mathcal{M}' :

$$F_2(a, b) : F(a) \otimes F(b) \rightarrow F(a \otimes b)$$

which are natural in a and b ,

- for the units e and e' , a morphism in \mathcal{M}' :

$$F_0 : e \rightarrow e'$$

making the diagrams commute:

$$\begin{array}{ccc} F(a) \otimes (F(b) \otimes F(c)) & \xrightarrow{\alpha'} & (F(a) \otimes F(b)) \otimes F(c) \\ \downarrow F(a) \otimes F_2(b, c) & & \downarrow F_2(a, b) \otimes F(c) \\ F(a) \otimes F(b \otimes c) & & F(a \otimes b) \otimes F(c) \\ \downarrow F_2(a, b \otimes c) & & \downarrow F_2(a \otimes b, c) \\ F(a \otimes (b \otimes c)) & \xrightarrow{F(\alpha)} & F((a \otimes b) \otimes c) \end{array} \quad (10)$$

$$\begin{array}{ccc} F(b) \otimes e' & \xrightarrow{\rho} & F(b) \\ \downarrow F(b) \otimes F_0 & & \uparrow F(\rho) \\ F(b) \otimes F(e) & \xrightarrow{F_2(b, e)} & F(b \otimes e) \end{array} \quad \begin{array}{ccc} e' \otimes F(b) & \xrightarrow{\lambda} & F(b) \\ \downarrow F_0 \otimes F(b) & & \uparrow F(\lambda) \\ F(e) \otimes F(b) & \xrightarrow{F_2(e, b)} & F(e \otimes b) \end{array} \quad (11)$$

A monoidal functor f is said to be *strong* when F_0 and all $F_2(a, b)$ are isomorphisms, *strict* when F_0 and all $F_2(a, b)$ are identities.

5.4. Braided monoidal functors

If \mathcal{M} and \mathcal{M}' are braided monoidal categories, a braided monoidal functor is a monoidal functor $(F, F_2, F_0) : \mathcal{M} \rightarrow \mathcal{M}'$ which commutes with the braidings γ and γ' in the following sense:

$$\begin{array}{ccc} F(a) \otimes F(b) & \xrightarrow{\gamma'} & F(b) \otimes F(a) \\ F_2(a, b) \downarrow & & \downarrow F_2(b, a) \\ F(a \otimes b) & \xrightarrow{F(\gamma)} & F(b \otimes a) \end{array} \quad (12)$$

5.5. Monoidal natural transformations

A monoidal natural transformation $\theta : (F, F_2, F_0) \rightarrow (G, G_2, G_0) : \mathcal{M} \rightarrow \mathcal{M}'$ between two monoidal functors is a natural transformation between the underlying ordinary functors $\theta : F \rightarrow G$ making the diagrams

$$\begin{array}{ccc} F(a) \otimes F(b) & \xrightarrow{F_2(a, b)} & F(a \otimes b) \\ \theta_a \otimes \theta_b \downarrow & & \downarrow \theta_{a \otimes b} \\ G(a) \otimes G(b) & \xrightarrow{G_2(a, b)} & G(a \otimes b) \end{array} \quad \begin{array}{ccc} e' & \xrightarrow{F_0} & Fe \\ \parallel & & \downarrow \theta_e \\ e' & \xrightarrow{G_0} & Ge \end{array} \quad (13)$$

commute in \mathcal{M}' .